arXiv:1703.03287v1 [math.CA] 9 Mar 2017

A REMARK ON CERTAIN INTEGRAL OPERATORS OF FRACTIONAL TYPE

PABLO A. ROCHA

Abstract. For $m, n \in \mathbb{N}$, $1 < m \leq n$, we write $n = n_1 + ... + n_m$ where $\{n_1, ..., n_m\} \subset \mathbb{N}$. Let $A_1, ..., A_m$ be $n \times n$ singular real matrices such that

$$\bigoplus_{i=1}^{m} \bigcap_{1 \leq j \neq i \leq m} \mathcal{N}_j = \mathbb{R}^n,$$

where $\mathcal{N}_j = \{x : A_j x = 0\}$, $dim(\mathcal{N}_j) = n - n_j$ and $A_1 + ... + A_m$ is invertible. In this paper we study integral operators of the form

$$Tf(x) = \int_{\mathbb{R}^n} |x - A_1 y|^{-n_1 + \alpha_1} \cdots |x - A_m y|^{-n_m + \alpha_m} f(y)\, dy,$$

$n = n_1 + ... + n_m$, $\frac{\alpha_1}{n_1} = ... = \frac{\alpha_m}{n_m} = r$, $0 < r < 1$, and the matrices A_i's are as above. We obtain the $H^p(\mathbb{R}^n) - L^q(\mathbb{R}^n)$ boundedness of T for $0 < p < \frac{1}{r}$ and $\frac{1}{q} = \frac{1}{p} - r$.

1. Introduction

For $0 \leq \alpha < n$ and $m > 1$, $(m \in \mathbb{N})$, let $T_{\alpha,m}$ be the integral operator defined by

$$T_{\alpha,m} f(x) = \int_{\mathbb{R}^n} |x - A_1 y|^{-\alpha_1} \cdots |x - A_m y|^{-\alpha_m} f(y) dy, \tag{1}$$

where $\alpha_1, ..., \alpha_m$ are positive constants such that $\alpha_1 + ... + \alpha_m = n - \alpha$, and $A_1, ..., A_m$ are $n \times n$ invertible matrices such that $A_i \neq A_j$ if $i \neq j$. We observe that for the case $\alpha > 0$, $m = 1$, and $A_1 = I$, $T_{\alpha,1}$ is the Riesz potential I_α. Thus for $0 < \alpha < n$ the operator $T_{\alpha,m}$ is a kind of generalization of the Riesz potential. The case $\alpha = 0$ and $m > 1$ was studied under the additional assumption that $A_i - A_j$ are invertible if $i \neq j$. The behavior of this class of operators and generalizations of them on the spaces of functions $L^p(\mathbb{R}^n)$, $L^p(w)$, $H^p(\mathbb{R}^n)$ and $H^p(w)$ was studied in [1], [2], [4], [5], [7] and [8].

If $0 < \alpha < n$ and $m > 1$, then the operator $T_{\alpha,m}$ has the same behavior that the Riesz potential on $L^p(\mathbb{R}^n)$. Indeed

$$|T_{\alpha,m} f(x)| \leq C \sum_{j=1}^{m} \int_{\mathbb{R}^n} |A_j^{-1} x - y|^{\alpha - n} |f(y)| dy = C \sum_{j=1}^{m} I_\alpha(|f|)(A_j^{-1} x),$$

for all $x \in \mathbb{R}^n$, this pointwise inequality implies that $T_{\alpha,m}$ is a bounded operator from $L^p(\mathbb{R}^n)$ into $L^q(\mathbb{R}^n)$ for $1 < p < \frac{n}{\alpha}$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, and it is of type weak $(1, n/n - \alpha)$.

It is well known that the Riesz potential I_α is bounded from $H^p(\mathbb{R}^n)$ into $H^q(\mathbb{R}^n)$ for $0 < p \leq 1$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ (see [11] and [3]). In [8], the author jointly with M. Urciuolo proved the $H^p(\mathbb{R}^n) - L^q(\mathbb{R}^n)$ boundedness of the operator $T_{\alpha,m}$ and we also showed that the $H^p(\mathbb{R}) - H^q(\mathbb{R})$ boundedness does not hold for $0 < p \leq \frac{1}{1+\alpha}$,

Key words and phrases: Integral operators, Hardy spaces.
2.010 Math. Subject Classification: 42B20, 42B30.
Partially supported by CONICET.

$\frac{1}{q} = \frac{1}{p} - \alpha$ and $T_{\alpha,m}$ with $0 \leq \alpha < 1$, $m = 2$, $A_1 = 1$, and $A_2 = -1$. This is a significant difference with respect to the case $0 < \alpha < 1$, $n = m = 1$, and $A_1 = 1$.

In this note we will prove that if we consider certain singular matrices in (1), then such an operator is still bounded from H^p into L^q. More precisely, for $m, n \in \mathbb{N}$, $1 < m \leq n$, we write $n = n_1 + ... + n_m$ where $\{n_1, ..., n_m\} \subset \mathbb{N}$, we also consider $n \times n$ singular real matrices $A_1, ..., A_m$ such that

$$\bigoplus_{i=1}^{m} \bigcap_{1 \leq j \neq i \leq m} \mathcal{N}_j = \mathbb{R}^n,$$

where $\mathcal{N}_j = \{x : A_j x = 0\}$, $dim(\mathcal{N}_j) = n - n_j$, $A_1 + ... + A_m$ is invertible. Given $0 < r < 1$ and $n_1, ..., n_m$ such that $n_1 + ... + n_m = n$, let $\alpha_1, ..., \alpha_m$ be positive constants such that $\frac{\alpha_1}{n_1} = ... = \frac{\alpha_m}{n_m} = r$, for such parameters we define the integral operator T_r by

$$T_r f(x) = \int_{\mathbb{R}^n} |x - A_1 y|^{-n_1+\alpha_1} \cdots |x - A_m y|^{-n_m+\alpha_m} f(y)\, dy, \tag{2}$$

where the matrices A_i's are as above.

We observe that the operator defined in (2) can be written as in (1) taking there the matrices A_i's singular. In fact, $T_r = T_{\beta,m}$ with $\beta_i = n_i - \alpha_i$ for each $i = 1, 2, ..., m$ and $\beta = nr$.

Our main result is the following

Theorem 1. *Let T_r be the integral operator defined in (2). If $0 < r < 1$, $0 < p < \frac{1}{r}$ and $\frac{1}{q} = \frac{1}{p} - r$, then T_r can be extended to an $H^p(\mathbb{R}^n) - L^q(\mathbb{R}^n)$ bounded operator.*

In Section 2 we state two auxiliary lemmas to get the main result in Section 3. We conclude this note with an example in Section 4.

Throughout this paper, c will denote a positive constant, not necessarily the same at each occurrence. The symbol $A \lesssim B$ stands for the inequality $A \leq cB$ for some constant c.

2. Preliminary results

Let K be a kernel in $\mathbb{R}^n \times \mathbb{R}^n$, we formally define the integral operator T_K by $T_K f(x) = \int_{\mathbb{R}^n} K(x, y) f(y) dy$.

We start with the following

Lemma 1. *Let $m, n \in \mathbb{N}$, with $1 < m \leq n$, and let $n_1, ..., n_m$ be natural numbers such that $n_1 + ... + n_m = n$. For each $i = 1, ..., m$ let K_i be kernels in $\mathbb{R}^{n_i} \times \mathbb{R}^{n_i}$ such that the operator T_{K_i} is bounded from $L^p(\mathbb{R}^{n_i})$ into $L^q(\mathbb{R}^{n_i})$ with $1 < p, q < \infty$, then the operator $T_{K_1 \otimes \cdots \otimes K_m}$ is bounded from $L^p(\mathbb{R}^n)$ into $L^q(\mathbb{R}^n)$.*

Proof. Since $\mathbb{R}^n = \mathbb{R}^{n_1} \times \cdots \times \mathbb{R}^{n_m}$ let $x = (x^1, ..., x^m) \in \mathbb{R}^{n_1} \times \cdots \times \mathbb{R}^{n_m}$, now the operator $T_{K_1 \otimes \cdots \otimes K_m}$ is given by

$$T_{K_1 \otimes \cdots \otimes K_m} f(x) = \int_{\mathbb{R}^{n_1} \times \cdots \times \mathbb{R}^{n_m}} K_1(x^1, y^1) \cdots K_m(x^m, y^m) f(y^1, ..., y^m)\, dy^1 ... dy^m.$$

Using that the kernels K_i define bounded operators for $1 \leq i \leq m$, the lemma follows from an iterative argument and the Minkowski's inequality for integrals.■

Lemma 2. *Let $m, n \in \mathbb{N}$, with $1 < m \leq n$, and let $n_1, ..., n_m$ be natural numbers such that $n_1 + ... + n_m = n$. If $A_1, ..., A_m$ are $n \times n$ singular real matrices such that*

$$\bigoplus_{i=1}^{m} \bigcap_{1 \leq j \neq i \leq m} \mathcal{N}_j = \mathbb{R}^n,$$

where $\mathcal{N}_j = \{x : A_j x = 0\}$, $dim(\mathcal{N}_j) = n - n_j$, and $A_1 + ... + A_m$ is invertible, then there exist two $n \times n$ invertible matrices B and C such that $B^{-1} A_j C$ is the canonical projection from $\mathbb{R}^n$ on $\{0\} \times \cdots \times \mathbb{R}^{n_j} \times \cdots \times \{0\}$ for each $j = 1, ..., m$.

Proof. It is easy to check that

$$\bigoplus_{i=1}^{m} \bigcap_{1 \leq j \neq i \leq m} \mathcal{N}_j = \mathbb{R}^n \Rightarrow \bigoplus_{1 \leq i \neq k \leq m} \bigcap_{1 \leq j \neq i \leq m} \mathcal{N}_j = \mathcal{N}_k$$

so

$$A_k \left(\bigcap_{1 \leq j \neq k \leq m} \mathcal{N}_j \right) = \mathcal{R}(A_k), \qquad k = 1, ..., m, \tag{3}$$

since $dim(\mathcal{N}_k) = n - n_k$ then $dim\left(\bigcap_{1 \leq j \neq k \leq m} \mathcal{N}_j\right) = dim(\mathcal{R}(A_k)) = n_k$. Let $\{\gamma_1^k, ..., \gamma_{n_k}^k\}$ be a basis of $\bigcap_{1 \leq j \neq k \leq m} \mathcal{N}_j$ thus $\{\gamma_1^1, ..., \gamma_{n_1}^1, ..., \gamma_1^m, ..., \gamma_{n_m}^m\}$ is a basis for $\mathbb{R}^n$. Let C be the $n \times n$ matrix which columns are the elements of the above basis. Since $A_1 + ... + A_m$ is invertible we have that $B = (A_1 + ... + A_m)C$ is invertible, so (3) gives that $B^{-1} A_j C$ is the canonical projection from $\mathbb{R}^n$ on $\{0\} \times \cdots \times \mathbb{R}^{n_j} \times \cdots \times \{0\}$ for each $j = 1, ..., m$. ■

3. The main result

Proof of the Theorem 1. We begin by obtaining the $L^p - L^q$ boundedness of the operator T_r for $1 < p < \frac{1}{r}$ and $\frac{1}{q} = \frac{1}{p} - r$, and then with this result we will prove the $H^p - L^q$ boundedness of T_r for $0 < p \leq 1$ and $\frac{1}{q} = \frac{1}{p} - r$.

$L^p - L^q$ boundedness. If A is a $n \times n$ invertible matrix we put $f_A(x) = f(A^{-1}x)$. Let B and C be the matrices given by Lemma 2, then

$$[T_r(f_C)]_{B^{-1}}(x) =$$

$$\int_{\mathbb{R}^n} |(Bx - A_1 y)|^{-n_1 + \alpha_1} \cdots |(Bx - A_m y)|^{-n_m + \alpha_m} f(C^{-1}y)\, dy =$$

$$|det(C)| \int_{\mathbb{R}^n} \left|B(x - B^{-1}A_1 C y)\right|^{-n_1 + \alpha_1} \cdots \left|B(x - B^{-1}A_m C y)\right|^{-n_m + \alpha_m} f(y)\, dy.$$

Since B is invertible, then there exists a positive constant c such that $c|x| \leq |Bx|$ for all $x \in \mathbb{R}^n$. Thus

$$|[T_r(f_C)]_{B^{-1}}(x)| \leq c \int_{\mathbb{R}^n} \left|x - B^{-1}A_1 C y\right|^{-n_1 + \alpha_1} \cdots \left|x - B^{-1}A_m C y\right|^{-n_m + \alpha_m} |f(y)|\, dy$$

$$\leq c \int_{\mathbb{R}^{n_1} \times ... \times \mathbb{R}^{n_m}} \left|x^1 - y^1\right|^{-n_1 + \alpha_1} \cdots |x^m - y^m|^{-n_m + \alpha_m} |f(y^1, ..., y^m)|\, dy^1 ... dy^m,$$

the second inequality it follows from Lemma 2 and from that $|x^j - y^j| \leq |x - P_j y|$, where $P_j = B^{-1}A_j C$ is the canonical projection from $\mathbb{R}^n$ on $\{0\} \times \cdots \times \mathbb{R}^{n_j} \times \cdots \times \{0\}$. Since $\gamma(\alpha_j)^{-1} |x^j - y^j|^{-n_j + \alpha_j}$, for an appropriate constant $\gamma(\alpha_j)$ (see [9], p. 117), is the kernel of the Riesz potential on $\mathbb{R}^{n_j}$, then Theorem 1 (in [9], p. 119) and

Lemma 1 give the $L^p - L^q$ boundedness of the operator T_r for $1 < p < \frac{1}{r}$ and $\frac{1}{q} = \frac{1}{p} - r$.

$H^p - L^q$ boundedness. Let $0 < p \leq 1$. We recall that a p-atom is a measurable function a supported on a ball B of $\mathbb{R}^n$ satisfying

a) $\|a\|_\infty \leq |B|^{-\frac{1}{p}}$

b) $\int y^\beta a(y)dy = 0$ for every multiindex β with $|\beta| \leq \lfloor n(p^{-1} - 1) \rfloor$, ($\lfloor s \rfloor$ denotes the integer part of s).

Let $0 < p \leq 1 < p_0 < \frac{1}{r}$, $0 < r < 1$, and $\frac{1}{q} = \frac{1}{p} - r$. Given $f \in H^p(\mathbb{R}^n) \cap L^{p_0}(\mathbb{R}^n)$, from Theorem 2, p. 107, in [10] we have that there exist a sequence of nonnegative numbers $\{\lambda_j\}_{j=1}^\infty$, a sequence of balls $B_j = B(z_j, \delta_j)$ centered at z_j with radius δ_j and p - atoms a_j supported on B_j, satisfying

$$\sum_{j \in \mathbb{N}} |\lambda_j|^p \leq c \|f\|_{H^p}^p , \tag{4}$$

such that f can be decomposed as $f = \sum\limits_{j \in \mathbb{N}} \lambda_j a_j$, where the convergence is in H^p and in L^{p_0} (for the converge in L^{p_0} see Theorem 5 in [6]). So the $H^p - L^q$ boundedness of T_r will be proved if we show that there exists $c > 0$ such that

$$\|T_r a_j\|_{L^q} \leq c \tag{5}$$

with c independent of the p-atom a_j. Indeed, since $f = \sum_{j=1}^\infty \lambda_j a_j$ in L^{p_0} and T_r is a $L^{p_0} - L^{\frac{p_0}{1 - r p_0}}$ bounded operator, we have that $|T_r f(x)| \leq \sum_{j=1}^\infty \lambda_j |T_r a_j(x)|$ a.e.x, this pointwise estimate, the inequality in (5), joint to the inequality

$$\left(\sum_{j \in \mathbb{N}} |\lambda_j|^{\min\{1,q\}} \right)^{\frac{1}{\min\{1,q\}}} \leq \left(\sum_{j \in \mathbb{N}} |\lambda_j|^p \right)^{\frac{1}{p}}$$

and (4) we obtain $\|T_r f\|_q \leq c \|f\|_{H^p}$, for all $f \in H^p(\mathbb{R}^n) \cap L^{p_0}(\mathbb{R}^n)$, so the theorem follows from the density of $H^p(\mathbb{R}^n) \cap L^{p_0}(\mathbb{R}^n)$ in $H^p(\mathbb{R}^n)$.

We will prove the estimate in (5). We define $D = \max_{1 \leq i \leq m} \max_{|y|=1} |A_i(y)|$. Let a_j be an p - atom supported on a ball $B_j = B(z_j, \delta_j)$, for each $1 \leq i \leq m$ let $B_{ji}^* = B(A_i z_j, 4D\delta_j)$. Since T_r is bounded from $L^{p_0}(\mathbb{R}^n)$ into $L^{q_0}(\mathbb{R}^n)$ for $\frac{1}{q_0} = \frac{1}{p_0} - r$, $1 < p_0 < \frac{1}{r}$, the Hölder inequality gives

$$\int\limits_{\bigcup\limits_{1 \leq i \leq m} B_{ji}^*} |T_r a_j(x)|^q \, dx \leq \sum_{1 \leq i \leq m} \int\limits_{B_{ji}^*} |T_r a_j(x)|^q \, dx$$

$$\leq c \sum_{1 \leq i \leq m} \left| B_{ji}^* \right|^{1 - \frac{q}{q_0}} \|T_r a_j\|_{q_0}^q \leq c \delta_j^{n - \frac{nq}{q_0}} \|a_j\|_{p_0}^q$$

$$\leq c \delta_j^{n - \frac{nq}{q_0}} \left(\int\limits_{B_j} |a_j|^{p_0} \right)^{\frac{q}{p_0}} \leq c \delta_j^{n - \frac{nq}{q_0}} \delta_j^{-\frac{nq}{p}} \delta_j^{\frac{nq}{p_0}} = c.$$

We denote $k(x,y) = |x - A_1 y|^{-n_1 + \alpha_1} \ldots |x - A_m y|^{-n_m + \alpha_m}$ and we put $N - 1 = \lfloor n(p^{-1} - 1) \rfloor$. In view of the moment condition of a_j we have

$$T_r a_j(x) = \int\limits_{B_j} k(x,y) a_j(y) dy = \int\limits_{B_j} \left(k(x,y) - q_{N,j}(x,y) \right) a_j(y) dy, \tag{6}$$

where $q_{N,j}$ is the degree $N-1$ Taylor polynomial of the function $y \to k(x,y)$ expanded around z_j. By the standard estimate of the remainder term of the taylor expansion, there exists ξ between y and z_j such that

$$|k(x,y)-q_{N,j}(x,y)| \lesssim |y-z_j|^N \sum_{k_1+\ldots+k_n=N} \left| \frac{\partial^N}{\partial y_1^{k_1}\ldots\partial y_n^{k_n}} k(x,\xi) \right|$$

$$\lesssim |y-z_j|^N \left(\prod_{i=1}^m |x-A_i\xi|^{-n_i+\alpha_i} \right) \left(\sum_{l=1}^m |x-A_l\xi|^{-1} \right)^N .$$

Now we decompose $\mathbb{R}^n = \bigcup_{i=1}^m B_{ji}^* \cup R_j$, where $R_j = \left(\bigcup_{i=1}^m B_{ji}^*\right)^c$, at the same time we decompose $R_j = \bigcup_{k=1}^m R_{jk}$ with

$$R_{jk} = \{x \in R_j : |x-A_kz_j| \leq |x-A_iz_j| \ for\ all\ i \neq k\}.$$

If $x \in R_j$ then $|x-A_iz_j| \geq 4D\delta_j$, for all $i=1,\ldots,m$, since $\xi \in B_j$ it follows that $|A_iz_j - A_i\xi| \leq D\delta_j \leq \frac{1}{4}|x-A_iz_j|$ so

$$|x-A_i\xi| = |x-A_iz_j+A_iz_j-A_i\xi| \geq |x-A_iz_j| - |A_iz_j-A_i\xi| \geq \frac{3}{4}|x-A_iz_j|.$$

If $x \in R_j$, then $x \in R_{jk}$ for some k, and since $\sum_{i=1}^m(-n_i+\alpha_i) = -n(1-r)$ we obtain

$$|k(x,y)-q_{N,j}(x,y)| \lesssim |y-z_j|^N \left(\prod_{i=1}^m |x-A_iz_j|^{-n_i+\alpha_i} \right) \left(\sum_{l=1}^m |x-A_lz_j|^{-1} \right)^N ,$$

$$\lesssim |y-z_j|^N |x-A_kz_j|^{-n(1-r)-N}, \quad if\ x \in R_{jk}\ \ and\ \ y \in B_j.$$

This inequality allow us to conclude that

$$\int_{R_j} \left| \int_{B_j} K(x,y)a_j(y)dy \right|^q dx = \int_{R_j} \left| \int_{B_j} [K(x,y)-q_{N,j}(x,y)]\, a_j(y)dy \right|^q dx$$

$$\lesssim \sum_{k=1}^m \int_{R_{jk}} \left(\int_{B_j} |y-z_j|^N |x-A_kz_j|^{-n(1-r)-N} |a_j(y)|\, dy \right)^q dx$$

$$\lesssim \left(\int_{B_j} |y-z_j|^N |a_j(y)|\, dy \right)^q \sum_{k=1}^m \int_{\left(B_{jk}^*\right)^c} |x-A_kz_j|^{-n(1-r)q-Nq}\, dx$$

$$\lesssim \delta_j^{qN-n\frac{q}{p}+nq} \int_{4D\delta_j}^{\infty} t^{-q(n(1-r)+N)+n-1} dt \leq c$$

with c independent of the $p-$atom a_j since $-q\left(n(1-r)+N\right)+n<0$.■

4. An example

For $n = m = 3$, $n_1 = n_2 = n_3 = 1$ we consider the following 3×3 singular matrices

$$A_1 = \begin{pmatrix} 4 & 4 & -1 \\ 0 & 0 & 0 \\ -4 & -4 & 1 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 1 & -1 & 0 \\ -2 & 2 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad A_3 = \begin{pmatrix} 1 & 0 & -1 \\ -3 & 0 & 3 \\ -1 & 0 & 1 \end{pmatrix}.$$

It is clear that

$$A_1 + A_2 + A_3 = \begin{pmatrix} 6 & 3 & -2 \\ -5 & 2 & 3 \\ -5 & -4 & 2 \end{pmatrix}$$

is invertible. For each $j = 1, 2, 3$, let $\mathcal{N}_j = \{x \in \mathbb{R}^3 : A_j x = 0\}$. A computation gives

$$\mathcal{N}_1 = \langle (1,0,4), (0,1,4) \rangle, \ \mathcal{N}_2 = \langle (1,1,0), (0,0,1) \rangle, \ \mathcal{N}_3 = \langle (1,0,1), (0,1,0) \rangle,$$

one can check that

$$\mathcal{N}_1 \cap \mathcal{N}_2 = \langle (1,1,8) \rangle, \ \mathcal{N}_1 \cap \mathcal{N}_3 = \langle (4,-3,4) \rangle, \ \textit{and} \ \mathcal{N}_2 \cap \mathcal{N}_3 = \langle (1,1,1) \rangle.$$

We observe that $\mathcal{N}_1 \cap \mathcal{N}_2 \oplus \mathcal{N}_1 \cap \mathcal{N}_3 \oplus \mathcal{N}_2 \cap \mathcal{N}_3 = \mathbb{R}^3$. As in the proof of Lemma 2, we define the matrices C and B by

$$C = \begin{pmatrix} 1 & 4 & 1 \\ 1 & -3 & 1 \\ 1 & 4 & 8 \end{pmatrix}, \ B = (A_1 + A_2 + A_3)\, C = \begin{pmatrix} 7 & 7 & -7 \\ 0 & -14 & 21 \\ -7 & 0 & 7 \end{pmatrix},$$

both matrices are invertibles with

$$B^{-1} = \begin{pmatrix} \frac{2}{21} & \frac{1}{21} & -\frac{1}{21} \\ \\ \frac{1}{7} & 0 & \frac{1}{7} \\ \\ \frac{2}{21} & \frac{1}{21} & \frac{2}{21} \end{pmatrix}.$$

Now it is easy to check that

$$B^{-1}A_1C = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \ B^{-1}A_2C = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \ B^{-1}A_3C = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

So, from Theorem 1, it follows that the operator T_r defined by

$$T_r f(x) = \int_{\mathbb{R}^3} |x - A_1 y|^{-1+r} |x - A_2 y|^{-1+r} |x - A_3 y|^{-1+r} f(y) dy,$$

with $0 < r < 1$, is a bounded operator from $H^p(\mathbb{R}^3)$ into $L^q(\mathbb{R}^3)$ for $0 < p < 1/r$ and $\frac{1}{q} = \frac{1}{p} - r$.

References

[1] T. GODOY, M. URCIUOLO, About the L^p boundedness of some integral operators, Revista de la UMA 38, 192-195, (1993).

[2] T. GODOY and M. URCIUOLO, On certain integral operators of fractional type, Acta Mathematica Hungar. (82), 99-105, (1999).

[3] S. KRANTZ, Fractional integration on Hardy spaces, Studia Mathematica, vol 73 (2), 87-94, (1982).

[4] M. S. RIVEROS and M. URCIUOLO, Weighted inequalities for integral operators with some homogeneous kernels, Czechoslovak Math. J., 55 (130), (2005), 423-432.

[5] M. S. RIVEROS and M. URCIUOLO, Weighted $H^p - L^q$ boundedness of integral operators with rough kernels. Preprint.

[6] P. ROCHA, A note on Hardy spaces and bounded linear operators . To appear in Georgian Math. Journal.
[7] P. ROCHA and M. URCIUOLO, On the Hp-Lp boundedness of some integral operators. Georgian Math. Journal, Vol 18, (4), 801 - 808, (2011).
[8] P. ROCHA and M. URCIUOLO, On the Hp-Lq boundedness of some fractional integral operators. Czechoslovak Math. Journal, 62 (137), 625 - 635, (2012).
[9] E. STEIN, Singular Integrals and Differentiability Properties of Functions, Princeton Univ. Press, 1970.
[10] E. STEIN, Harmonic Analysis, Real Variable Methods, Orthogonality and Oscillatory Integrals, Princeton Univ. Press, 1993.
[11] M. H. TAIBLESON and G. WEISS, The molecular characterization of certain Hardy spaces, Asterisque 77, 67-149, (1980).

Universidad Nacional del Sur, INMABB (Conicet), Bahía Blanca, 8000 Buenos Aires, Argentina

E-mail address: pablo.rocha@uns.edu.ar